

MONTE CARLO SIMULATION OF THE THREE DIMENSIONAL THIRRING MODEL

Luigi Del Debbio and Simon Hands

*Department of Physics
 University of Wales, Swansea,
 Singleton Park,
 Swansea SA2 8PP, U.K.*

Abstract

We study the Thirring model in three spacetime dimensions, by means of Monte Carlo simulation on lattice sizes 8^3 and 12^3 , for numbers of fermion flavors $N_f = 2, 4, 6$. For sufficiently strong interaction strength, we find that spontaneous chiral symmetry breaking occurs for $N_f = 2, 4$, in accordance with the predictions of the Schwinger-Dyson approach. The phase transitions which occur are continuous and with critical scaling behaviour depending on N_f . For $N_f = 6$ our results are preliminary, and no firm conclusions about the existence or otherwise of chiral symmetry breaking are possible.

PACS numbers: 11.10.Kk, 11.15.Ha, 11.30.Rd

Keywords: Model field theory, $1/N$ expansion, Schwinger-Dyson equation, chiral symmetry breaking, lattice simulation, dynamical fermions.

The three dimensional Thirring model is a field theory of relativistic fermions interacting via a contact term between conserved vector currents. Its Lagrangian is written

$$\mathcal{L} = \bar{\psi}_i(\not{\partial} + m)\psi_i + \frac{g^2}{2N_f}(\bar{\psi}_i\gamma_\mu\psi_i)^2, \quad (1)$$

where $\psi_i, \bar{\psi}_i$ are four-component spinors, m is a bare, parity-conserving mass, and the index i runs over N_f distinct fermion species. Since the coupling g^2 has mass dimension -1 , naive power-counting suggests that the model (1) is non-renormalisable. However, as has been suspected for many years [1,2], an expansion in powers of $1/N_f$, rather than g^2 , is exactly renormalisable. At leading order in $1/N_f$, interaction between vector currents is dominated by exchange of a fermion - anti-fermion bound state described in terms of a chain of vacuum-polarisation “bubbles”. In the ultra-violet limit the interaction is thus transformed from a momentum-independent contact term to a softer $A/(N_fk)$ behaviour, where A is a numerical constant independent of g : this asymptotic behaviour can be used to evaluate divergent graphs at higher orders in $1/N_f$, eg. in [3], where renormalisability of the massless model is explicitly demonstrated to $O(1/N_f)$.

The property of renormalisability signals that the model’s $1/N_f$ expansion exhibits a UV-stable fixed point of the renormalisation group, the continuum limit being taken in the limit $g^2\Lambda \rightarrow \infty$, where Λ is a UV cutoff. RG fixed points have also been observed in other three-dimensional four-fermi models [4,5]. The distinctive feature of the Thirring model is that for $d < 4$ the vacuum polarisation is UV-finite so long as the regularisation respects current conservation (eg. Pauli-Villars). This means that there is no need to fine-tune g^2 to a critical value: a continuum limit may be taken for any value of the dimensionless parameter mg^2 (at least to leading order in $1/N_f$ [3]), the theory thus obtained having a variable ratio of, say, physical fermion mass to vector bound state mass. In the RG sense the interaction $(\bar{\psi}\gamma_\mu\psi)^2$ is a *marginal* operator for $2 < d < 4$, whereas, say, the interaction $(\bar{\psi}\psi)^2$ in the Gross-Neveu model is *relevant* [6].

Another possibility raised by the $1/N_f$ expansion is the equivalence of the Thirring

model in the strong-coupling limit $mg^2 \rightarrow \infty$, in which the vector particle becomes massless, with the infra-red limiting behaviour of QED in three spacetime dimensions. In massless large- N_f QED₃ vacuum polarisation screens one-photon exchange to the extent that the $1/k^2$ interaction is again transformed to $1/k$ [7]. The $O(1/N_f)$ corrections to the models evaluated in respectively UV (Thirring) or IR (QED) limits appear to coincide [3,8].

The $1/N_f$ expansion may not, however, describe the true behaviour of the model, particularly for small N_f . For instance, spontaneous chiral symmetry breaking, signalled by a vacuum condensate $\langle \bar{\psi}\psi \rangle$, is forbidden to all orders in $1/N_f$, and yet may be predicted by a self-consistent approach such as solution of the Schwinger-Dyson equations [2,9,10]. In this approach a non-trivial solution for the dressed fermion propagator $S(p) = (A(p)i\not{p} + \Sigma(p))^{-1}$ is sought, ie. one in which the self-energy $\Sigma(p)$ and hence $\langle \bar{\psi}\psi \rangle$ are non-vanishing in the chiral limit $m \rightarrow 0$. Unfortunately, the SD equations can only be solved by truncating them in a somewhat arbitrary fashion. The usual approximation [2,10] is to assume that the vector propagator is given by its leading-order form for $m = 0$ in the $1/N_f$ expansion, viz.

$$D_{\mu\nu}(k) = \left(\delta_{\mu\nu} - \frac{k_\mu k_\nu}{k^2} \right) \left(1 + \frac{g^2(k^2)^{\frac{1}{2}}}{8} \right)^{-1} + \frac{k_\mu k_\nu}{k^2}, \quad (2)$$

and that the fermion-vector vertex function is well-approximated by the bare vertex (the so-called “planar” or “ladder” approximation):

$$\Gamma_\mu(p, q) = -\frac{ig}{\sqrt{N_f}} \gamma_\mu. \quad (3)$$

The longitudinal part of $D_{\mu\nu}$ raises potential ambiguities: the most systematic treatment has been given by Itoh *et al* [10], who note the equivalence of the Thirring model with a gauge-fixed form of a fermion-scalar model possessing a local gauge symmetry and then use a non-local gauge-fixing condition to find a gauge in which the “wavefunction renormalisation” $A(p) \equiv 1$. This simplification enables the SD equations to be exactly

solved in the limit $g^2 \rightarrow \infty$, with the result that a non-trivial solution for $\Sigma(p)$ exists for

$$N_f < N_{fc} = \frac{128}{3\pi^2} \simeq 4.32. \quad (4)$$

Moreover since the integral equations require the introduction of a UV cutoff Λ , a feature of this solution is that the induced physical mass scale μ depends on N_f in an essentially singular way:

$$\frac{\mu}{\Lambda} \propto \exp\left(-\frac{2\pi}{\sqrt{\frac{N_{fc}}{N_f}-1}}\right); \quad \langle\bar{\psi}\psi\rangle \propto \Lambda^{\frac{1}{2}}\mu^{\frac{3}{2}} \propto \exp\left(-\frac{3\pi}{\sqrt{\frac{N_{fc}}{N_f}-1}}\right). \quad (5)$$

This implies that a continuum limit only exists as $N_f \rightarrow N_{fc}$, the scenario being very similar to that proposed by Miranskii and collaborators for strongly-coupled QED₄ [11]. Unfortunately no analytic solution exists for $g^2 < \infty$; however using different techniques Kondo [12] has argued that a critical line $N_{fc}(g^2)$ exists in the (g^2, N_f) plane, which is a smooth invertible function. Therefore for integer $N_f < N_{fc}$ one might expect a critical scaling behaviour

$$\langle\bar{\psi}\psi\rangle \propto \exp\left(-\frac{a}{\sqrt{\frac{g_c^2}{g^2}-1}}\right), \quad (6)$$

corresponding to a symmetry restoring transition at some critical point $g^2 = g_c^2$. Presumably in this scenario the Thirring interaction has become relevant: there may exist a novel strongly-coupled continuum limit at the critical point not described by the $1/N_f$ expansion.

There are good reasons to be cautious of this picture, however. Using a different sequence of truncations Hong and Park have found chiral symmetry breaking for all N_f [9], with

$$\frac{1}{g_c^2} \propto \exp\left(-\frac{N_f\pi^2}{16}\right), \quad (7)$$

a result which is non-analytic in $1/N_f$. Moreover in the limit $g^2 \rightarrow \infty$ the system of SD equations obtained are very similar to those of large- N_f QED₃, in which case studies

beyond the planar approximation, using improved ansätze for the vertex Γ_μ , suggest that the condition $A(p) \equiv 1$ is unphysical, and that chiral symmetry is spontaneously broken for all N_f [13]. In the current context this would imply $g_c^2 < \infty$ for all N_f .

For these reasons we consider a numerical study of the lattice-regularised model to be timely. If, as suggested above, the Thirring model lies in the same universality class as QED_3 , then a numerical study may shed light on the value of N_{fc} for this model [7]; previous lattice studies [14] have been plagued by large finite volume effects due to the slow fall-off of the photon propagator $\propto 1/x$. The corresponding propagator in the Thirring model falls as $1/x^2$, so sensible results may emerge on smaller systems. A second motivation is the possible existence of a novel continuum limit: since vacuum polarisation corrections to the vector propagator are finite to all orders in $1/N_f$ there should be no competing effects of charge screening, which obscures the issue in QED_4 .

The lattice action we have used is as follows:

$$\begin{aligned}
S &= \frac{1}{2} \sum_{x\mu i} \bar{\chi}_i(x) \eta_\mu(x) (\chi_i(x + \hat{\mu}) - \chi_i(x - \hat{\mu})) + m \sum_{xi} \bar{\chi}_i(x) \chi_i(x) \\
&\quad + \frac{g^2}{2N} \sum_{x\mu ij} \bar{\chi}_i(x) \chi_i(x + \hat{\mu}) \bar{\chi}_j(x + \hat{\mu}) \chi_j(x) \\
&= \frac{1}{2} \sum_{x\mu i} \bar{\chi}_i(x) \eta_\mu(x) (1 + iA_\mu(x)) \chi_i(x + \hat{\mu}) + \text{h.c.} + m \sum_{xi} \bar{\chi}_i(x) \chi_i(x) + \frac{N}{4g^2} \sum_{x\mu} A_\mu^2(x), \\
&= \frac{1}{2} \sum_{xy\mu i} \bar{\chi}_i(x) M_{(A,m)}(x, y) \chi_i(y) + \frac{N}{4g^2} \sum_{x\mu} A_\mu^2(x),
\end{aligned} \tag{8}$$

where $\chi, \bar{\chi}$ are staggered fermion fields, η_μ the Kawamoto-Smit phases, m is the bare mass, the flavor index i runs from 1 to N , and we have introduced $M_{(A,m)}$ for the fermionic bilinear, which depends on both the auxiliary field and the mass. The second form of the action is the one actually simulated: the equivalence of the two forms follows from Gaussian integration over the real-valued auxiliary field A_μ defined on the lattice links (for $N = 1$ there is an alternative formulation in terms of a compact complex-valued auxiliary

[15,16]). The vector-like interaction of the action allows the introduction of a checker-board, which in turn enables simulation of the system for any N . In three Euclidean dimensions staggered fermions describe two continuum species of four-component fermions, with a parity-conserving mass term [17]. Hence the number of physical flavors $N_f = 2N$. An interesting feature of the lattice formulation (8) is that the interaction current is not exactly conserved. The conserved current in lattice gauge theory incorporates the gauge connection $\exp(iA_\mu)$. This means that at leading order in $1/N$ the vector propagator receives an extra contribution from vacuum polarisation, essentially due to the absence of the diagram of Fig. 1. The effect can be absorbed into a redefinition of the coupling:

$$g_R^2 = \frac{g^2}{1 - g^2 J(m)}, \quad (9)$$

where $J(m)$ is the value of the integral depicted in Fig. 1. The physics described by continuum $1/N_f$ perturbation theory occurs for the range of couplings $g_R^2 \in [0, \infty)$, ie. for $g^2 \in [0, g_{lim}^2)$; to leading order in $1/N$

$$\frac{1}{g_{lim}^2} = J(m); \quad \text{with} \quad J(0) = \frac{2}{3}. \quad (10)$$

We therefore expect to see some kind of discontinuous behaviour in our simulations for small values of $1/g^2$.

In this letter, we aim to clarify the chiral symmetry breaking pattern by studying the chiral condensate:

$$\langle \bar{\psi} \psi \rangle = \frac{1}{V} \text{Tr} \left(M_{(A,m)}^{-1} \right)$$

which, in the limit $m \rightarrow 0$, is an order parameter for the spontaneous symmetry breaking. We performed simulations on 8^3 and 12^3 lattices for $N_f = 2, 4, 6$, using the hybrid Monte Carlo algorithm. Bare mass values m ranged from 0.4 down to 0.02, with most attention paid to the range 0.05 – 0.02. For each mass and coupling we performed roughly 500 HMC trajectories, the trajectory length being drawn from a Poisson distribution with mean 0.9.

The condensate $\langle \bar{\psi}\psi \rangle$ was measured with a stochastic estimator every few trajectories. To maintain reasonable acceptance rates the timestep varied from 0.15 on 8^3 at $m = 0.4$ down to 0.022 on 12^3 at $m = 0.02$. We found that considerably more work was needed to perform matrix inversion in this model than for the Gross-Neveu simulations described in [5]. Another difference is that in this case since the critical region of interest occurs at successively stronger couplings as N_f is raised, the *CPU* required also grows with N_f , despite the $1/N_f$ suppression of quantum corrections.

In Fig. 2, we plot $\sigma \equiv \langle \bar{\psi}\psi \rangle$ vs. $1/g^2$ for the three values of N_f studied, for $m = 0.10$ on a 8^3 lattice. The models with different N_f have apparently coincident condensates in the strong-coupling region $1/g^2 \leq 0.3$, but thereafter the $\langle \bar{\psi}\psi \rangle$ signals peak to maxima at distinct values of $1/g^2$ before falling away. It is tempting to associate the strong-coupling region with $g_R^2 < 0$ from the discussion following Eq. (9), although the correspondence with the value of g_{lim}^2 predicted in Eq. (10) is not good. It may well be that the value of the diagram of Fig. 1 is considerably altered in a chirally broken vacuum.

As stated above, in order to study spontaneous chiral symmetry breaking, one has to monitor the value of $\langle \bar{\psi}\psi \rangle$ as $m \rightarrow 0$. Our results for the chiral condensate for $N_f = 2$ are reported in Fig. 3 for different values of the bare mass. A naive extrapolation to the chiral limit from the lattice data at fixed $1/g^2$ is probably unreliable in the range of bare masses we have explored. In order to determine the critical point, we need to perform a global fit of our data incorporating many values of m and $1/g^2$. Therefore, we have to use an equation of state (EOS) relating the external symmetry breaking parameter m to the response of the system $\langle \bar{\psi}\psi \rangle$ and the coupling $1/g^2$ [16,18].

A generic EOS, inspired by the critical behaviour of spin systems, can be written in terms of the scaled variables:

$$m\langle \bar{\psi}\psi \rangle^{-\delta} = \mathcal{F} \left(\Delta(1/g^2) \langle \bar{\psi}\psi \rangle^{1/\beta} \right), \quad (11)$$

where $\Delta(1/g^2) = 1/g_c^2 - 1/g^2$ is the reduced coupling. At $g = g_c$, Eq. (11) is the usual

scaling relation:

$$\langle \bar{\psi}\psi \rangle \sim m^{1/\delta},$$

while a Taylor expansion for small $\Delta(1/g^2)$ yields:

$$m = B\langle \bar{\psi}\psi \rangle^\delta + A\Delta(1/g^2)\langle \bar{\psi}\psi \rangle^{\delta-1/\beta} + \dots \quad (12)$$

where $A, B = \mathcal{F}'(0), \mathcal{F}(0)$ respectively. At this stage, one sees that, for vanishing m , Eq. (12) is simply the definition of the critical exponent β :

$$\langle \bar{\psi}\psi \rangle \sim (1/g_c^2 - 1/g^2)^\beta,$$

and that there are no logarithmic corrections, since these should only appear in 4-d [18]. If the critical behaviour is described by mean-field theory, we expect $\delta = 3$ and $\beta = 1/2$, yielding:

$$\langle \bar{\psi}\psi \rangle^2 = \kappa_1 \frac{m}{\langle \bar{\psi}\psi \rangle} + \kappa_2 \Delta(1/g^2), \quad (13)$$

which shows that $\langle \bar{\psi}\psi \rangle^2$ is a linear function of the ratio $m/\langle \bar{\psi}\psi \rangle$. Such a plot is known as Fisher plot. From Eq. (13) we see that a positive value of the intercept corresponds to a non-vanishing value of the chiral condensate for $m = 0$, while the intercept will be exactly zero at the critical coupling. In Fig. 4 and 5 we show the Fisher plots for $N_f = 2, 4$ respectively, where we can see at first glance an indication of chiral symmetry breaking, according to the criterion stated above. In order to get a more quantitative evidence, we have fitted our data using Eq. (12) and a simpler version of it based on the hypothesis that $\delta - 1/\beta = 1$ [19], which we will call respectively fit I and II in what follows. We should stress that in the absence of a systematic critical theory the forms I and II are used simply as effective descriptions of the data. The values of the fit parameters, their errors and the χ^2 are listed in Table 1. The number of values of the chiral condensate included in the fit is chosen in order to minimize the value of the reduced χ^2 . The results of fit II are shown in Figs. 4 and 5 and seem to describe the data quite well. The dashed line in Fig. 3 is the

curve one obtains using the results of fit II with $m = 0$. It shows clearly that, within the range of m we have explored, the chiral condensate is still far from its chiral limit value, thus providing a justification *a posteriori* for the impossibility of extrapolating the data naively.

There are a few conclusions one can draw from the numerical analysis that we would like to emphasize. First, for both values of N_f , we find clear evidence of chiral symmetry breaking at finite values of the coupling, as predicted by the Schwinger-Dyson approach. From Eq. (7) we get:

$$\frac{g_c^2(N_f = 2)}{g_c^2(N_f = 4)} = \exp(-\pi^2/8) = 0.291$$

which is not too far from the fitted value 0.342 ± 0.015 (using the data from fit II).

Secondly, although any claims to understand the details of the critical scaling must be premature, the fits strongly suggest that the models with $N_f = 2$ and 4 are described by distinct critical theories, in a sense on “either side” of the mean field theory. If we relax the requirement $\delta - 1/\beta = 1$ (which means using fit I instead of fit II) then the difference in the fitted values of δ becomes even more apparent. This is significant because similar EOS fits in QED₄ reveal no such differences between $N_f = 2$ and $N_f = 4$ [20].

Finally, we report some preliminary results for $N_f = 6$. Figure 6 shows $\langle \bar{\psi}\psi \rangle$ vs. m for two values of g . The $1/g^2 = 0.5$ data suggest a linear extrapolation to the chiral limit, yielding a small condensate equal to 0.013(4). For $1/g^2 = 0.4$ it is less clear how to make the extrapolation. Clearly in either case reliable data at much smaller mass values would be needed for confirmation or otherwise of chiral symmetry breaking: comparing data from different lattice sizes, we have found that finite size effects become more important as we go to larger N_f , which means that larger lattice sizes will probably be needed before we can proceed to a more quantitative study.

We conclude by briefly summarizing our results. We have shown by numerical simulations that spontaneous chiral symmetry breaking does occur in the Thirring model for

finite N_f , in contradiction with the $1/N_f$ perturbative expansion, but in agreement with the Schwinger-Dyson approach at least for $N_f = 2, 4$. In these two cases we were able to determine the critical coupling and the critical exponent δ by fitting to a plausible EOS, and the fits suggest that the two models have different critical behaviour. For $N_f = 6$, we were unable to find clear evidence in favour of symmetry breaking, but cannot yet exclude a non-vanishing condensate in the chiral limit. In the future we plan to investigate in more detail the theory for $N_f = 2, 4, 6$ at the critical point, focussing on critical exponents, the renormalized charge, and spectroscopy.

Acknowledgments

LDD is supported by an EC HMC Institutional Fellowship under contract No. ERBCH-BGCT930470, and SJH by a PPARC Advanced Fellowship. Some of the numerical work was performed on the Cray Y-MP at Rutherford–Appleton Laboratory under PPARC grant GR/J675.5. We have enjoyed discussing this project with Jiří Jersák, Kei-Ichi Kondo, Mike Pennington, Craig Roberts and Paolo Rossi.

References

- [1] G. Parisi, Nucl. Phys. **B100** (1975) 368;
S. Hikami and T. Muta, Prog. Theor. Phys. **57** (1977) 785;
Z. Yang, Texas preprint UTTG-40-90 (1990).
- [2] M. Gomes, R.S. Mendes, R.F. Ribeiro and A.J. da Silva, Phys. Rev. **D43** (1991) 3516.
- [3] S.J. Hands, Phys. Rev. **D51** (1995) 5816.
- [4] B. Rosenstein, B.J. Warr and S.H. Park, Phys. Rep. **205** (1991) 59.
- [5] S.J. Hands, A. Kocić and J.B. Kogut, Ann. Phys. **224** (1993) 29.
- [6] G. Gat, A. Kovner and B. Rosenstein, Nucl. Phys. **B5.5** (1992) 76.
- [7] R.D. Pisarski, Phys. Rev. **D29** (1984) 2423;
T.W. Appelquist, M. Bowick, D. Karabali and L.C.R. Wijewardhana, Phys. Rev. **D33** (1986) 3704.
- [8] D.Espriu, A. Palanques-Mestre, P. Pascual and R. Tarrach, Z. Phys. **C13** (1982) 153;
A. Palanques-Mestre and P. Pascual, Comm. Math. Phys. **95** (1984) 277.
- [9] D.K. Hong and S.H. Park, Phys. Rev. **D49** (1994) 5507.
- [10] T. Itoh, Y. Kim, M. Sugiura and K. Yamawaki, Prog. Theor. Phys. **93** (1995) 417.
- [11] P.I. Fomin, V.P. Gusynin, V.A. Miranskii and Yu.A. Sitenko, Riv. Nuovo Cimento **6** (1983) 1;
V.A. Miranskii, Nuovo Cimento **90A** (1985) 149.
- [12] K.-I. Kondo, Nucl. Phys. **B450** (1995) 251.
- [13] M.R. Pennington and S.P. Webb, Brookhaven preprint BNL-40886 (1988);
D. Atkinson, P.W. Johnson and M.R. Pennington, Brookhaven preprint BNL-41615 (1988);

- M.R.Pennington and D. Walsh, Phys. Lett. **B253** (1991) 246.
- [14] E. Dagotto, A. Kocić and J.B. Kogut, Nucl. Phys. **B334** (1990) 279;
S.J. Hands and J.B. Kogut, Nucl. Phys. **B5.5** (1990) 455.
- [15] S.P. Booth, R.D. Kenway and B.J. Pendleton, Phys. Lett. **B228** (1989) 115.
- [16] A. Ali Khan, M. Göckeler, R. Horsley, P.E.L. Rakow, G. Schierholz and H. Stüben,
Phys. Rev. **D51** (1995) 5.51.
- [17] C.J. Burden and A.N. Burkitt, Europhys. Lett. **3** (1987) 545.
- [18] M. Göckeler, R. Horsley, P.E.L. Rakow, G. Schierholz and R. Sommer, Nucl. Phys.
B371 (1992) 713.
- [19] E. Dagotto, S.J. Hands, A. Kocić, J.B. Kogut, Nucl. Phys. **B347** (1990) 217.
- [20] S.J. Hands, A. Kocić, J.B. Kogut, R.L. Renken, D.K. Sinclair, K.C. Wang, Nucl.
Phys. **B413** (1994) 503;
A. Kocić, J.B. Kogut, K.C. Wang, Nucl. Phys. **B398** (1993) 5.5.

param.	fit I	fit II
$N_f = 2$		
β_c	2.03(9)	1.94(4)
δ	2.32(23)	2.68(16)
β	0.71(9)	–
A	0.32(5)	0.37(1)
B	1.91(43)	2.86(53)
$\chi^2/\text{d.o.f}$	2.4	2.1
$N_f = 4$		
β_c	0.63(1)	0.66(1)
δ	3.67(28)	3.43(19)
β	0.38(4)	–
A	0.78(5)	0.73(2)
B	7.9(2.8)	6.4(1.5)
$\chi^2/\text{d.o.f}$	3.1	2.0

Table 1

Results from the fits

Figure Captions

Figure 1: Diagram contributing to coupling constant renormalisation at leading order in $1/N_f$.

Figure 2: chiral condensate σ vs. $1/g^2$ for $N = 1, 2, 3$, corresponding respectively to $N_f = 2, 4, 6$.

Figure 3: chiral condensate σ vs. $1/g^2$ for $N_f = 2$ and different values of the bare mass m .

Figure 4: Fisher plot for $N_f = 2$, from data at $\beta = 1.6(\triangle)$, $1.8(\triangleleft)$, $2.0(\nabla)$, $2.2(\triangleright)$, $2.4(+)$, $2.6(\times)$.

Figure 5: Fisher plot for $N_f = 4$, from data at $\beta = 0.5(\circ)$, $0.6(\blacksquare)$, $0.7(\diamond)$, $0.8(\triangle)$, $0.9(\triangleleft)$, $1.0(\nabla)$, $1.1(\triangleright)$, $1.2(+)$, $1.3(\times)$, $1.4(*)$.

Figure 6: chiral condensate σ vs. m for different values of the coupling (here $\beta \equiv 1/g^2$ and is not related to the critical exponent).




Figure 1

Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6

